## 5-Year Cause-Specific Survival<sup>a</sup> and Case Distribution by Race/Ethnicity

5-Year Cause-Specific Survival<sup>a</sup> and Case Distribution by Race/Ethnicity SEER 18 Registries, 2003-2009

Table 33.5

Cancer of the Colon and Rectum

	Both Sexes				Males				Females			
	Case Distribution		Survival		Case Distribution		Survival		Case Distribution		Survival	
Race/Ethnicity	Count	Percent <sup>b</sup>	Percent	SE	Count	Percent <sup>b</sup>	Percent	SE	Count	Percent <sup>b</sup>	Percent	SE
All Races	212,664	100.0	64.9	0.1	108,602	100.0	65.4	0.2	104,062	100.0	64.5	0.2
White	169,224	79.6	65.2	0.1	87,211	80.3	65.8	0.2	82,013	78.8	64.7	0.2
White Non-Hispanic	150,136	70.6	65.4	0.1	76,894	70.8	66.0	0.2	73,242	70.4	64.7	0.2
White Hispanic	19,088	9.0	64.2	0.4	10,317	9.5	64.1	0.6	8,771	8.4	64.3	0.6
Black	24,823	11.7	58.7	0.4	11,723	10.8	57.7	0.6	13,100	12.6	59.5	0.5
American Indian/Alaska Native <sup>c</sup>	1,034	0.5	62.7	1.8	518	0.5	61.8	2.6	516	0.5	63.4	2.4
Asian	14,797	7.0	69.9	0.5	7,563	7.0	70.0	0.6	7,234	7.0	69.8	0.6
Asian Indian, Pakistani	696	0.3	73.4	2.0	410	0.4	73.8	2.6	286	0.3	73.0	3.1
Chinese	3,463	1.6	70.9	0.9	1,736	1.6	71.6	1.3	1,727	1.7	70.3	1.3
Filipino	3,190	1.5	67.8	1.0	1,651	1.5	68.1	1.4	1,539	1.5	67.5	1.5
Japanese	2,942	1.4	69.3	1.0	1,473	1.4	68.2	1.4	1,469	1.4	70.3	1.4
Korean	1,440	0.7	69.5	1.4	709	0.7	69.4	2.1	731	0.7	69.7	2.0
Vietnamese	1,205	0.6	68.5	1.6	641	0.6	67.8	2.4	564	0.5	69.1	2.3
Other Asian <sup>d</sup>	1,861	0.9	72.5	1.2	943	0.9	73.6	1.7	918	0.9	71.3	1.8
Pacific Islander	918	0.4	59.8	2.0	541	0.5	57.6	2.7	377	0.4	63.0	3.1
Hawaiian	624	0.3	59.9	2.5	374	0.3	56.2	3.3	250	0.2	65.6	3.8
Other Pacific Islander <sup>e</sup>	294	0.1	59.5	3.6	167	0.2	60.9	4.7	127	0.1	57.9	5.3
$ ext{Hispanic}^{ ext{f}}$	19,833	9.3	64.2	0.4	10,680	9.8	64.3	0.6	9,153	8.8	64.2	0.6

Source: SEER 18 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana, New Jersey and Georgia excluding ATL/RG).

Five-Year Cause-Specific survival was calculated by actuarial method using diagnosis years 2003-2009 with follow-up through 2010.

b Percent of cases do not sum to 100 because All Races includes unknown race and other race which are not shown separately.

Survival estimates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

d Other Asians include Laotian, Hmong, Kampuchean, Thai, and Asians, Not Otherwise Specified (NOS).

Other Pacific Islanders include Micronesian, Chamorran, Guamanian, Polynesian, Tahitian, Samoan, Tongan, Melanesian, Fiji Islander, New Guinean, and Pacific Islander, NOS.

Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.

<sup>-</sup> Statistic not shown due to fewer than 25 cases during the time period.